

clooctadiene)rhodium(I)Tetrafluoroborate, (CAS #79255-71-3). The hexyne reaction was performed in a solution of 50% tetrahydrofuran with 5 mL total and 30 mG Wilkinson's Catalyst. Most of this volume does not contribute to signal since it is far from the magnetometer.

REFERENCES

- [0090] 1. Ernst, R. R., Bodenhausen, G., A. Wokaun, *Principles of Nuclear Magnetic Resonance in One and Two Dimensions*, Oxford University Press, New York (1987).
- [0091] 2. Slichter, C. P., *Principles of Magnetic Resonance, 3rd ed.* Springer-Verlag, New York (1990).
- [0092] 3. Appelt, S., Häsing, F. W., Kühn, H., Perlo, J., & Blümich, B. Mobile high resolution xenon nuclear magnetic resonance spectroscopy in the Earth's magnetic field. *Phys. Rev. Lett.* 94, 197601-197604 (2005).
- [0093] 4. McDermott, R. et al. Liquid-state NMR and scalar couplings in microtesla magnetic fields. *Science* 295, 2247-2249 (2002).
- [0094] 5. Ledbetter, M. P. et al. Optical detection of NMR J-spectra at zero magnetic field. *J. Magn. Res.* 199, 25-29, (2009).
- [0095] 6. Appelt, S., Kühn, H., Häsing, F. W., & Blümich, B. Chemical analysis by ultrahigh-resolution nuclear magnetic resonance in the Earth's magnetic field. *Nat. Phys.* 2, 105-109, (2006).
- [0096] 7. Budker, D. & Romalis, M. V. Optical Magnetometry. *Nature Physics* 3, 227-234 (2007).
- [0097] 8. Kominis, I. K., Kornack, T. W., Allred, J. C., & Romalis, M. V. A sub-femtoTesla multichannel atomic magnetometer, *Nature*. 422, 596-599 (2003).
- [0098] 9. Greenberg, Y. S. Application of superconducting quantum interference devices to nuclear magnetic resonance. *Reviews of Modern Physics* 70, 175-222 (1998).
- [0099] 10. Ledbetter, M. P. et al. Zero-field remote detection of NMR with a microfabricated atomic magnetometer. *Proc. Nat. Acad. Sci. (USA)* 105, 2286-2290, (2008).
- [0100] 11. Savukov, I. M. & Romalis, M. V. NMR detection with an atomic magnetometer. *Phys. Rev. Lett.* 94, 123001/1-4 (2005).
- [0101] 12. Xu, S. J. et al. Magnetic resonance imaging with an optical atomic magnetometer. *Proc. Nat. Acad. Sci. (USA)* 103, 12668-12671, (2006).
- [0102] 13. Savukov, I. M. et al. MRI with an atomic magnetometer suitable for practical imaging applications. *J. Magn. Res.* 199, 188-191 (2009).
- [0103] 14. Bowers, C. R. & Weitekamp, D. P. Transformation of symmetrization order to nuclear-spin magnetization by chemical-reaction and nuclear-magnetic-resonance. *Phys. Rev. Lett.* 57, 2645-2648 (1986).
- [0104] 15. Natterer, J. & Bargon, J. Parahydrogen induced polarization. *Prog in Nucl. Magn. Res. Spec.* 31, 293-315 (1997).
- [0105] 16. Bowers, C. R. Sensitivity enhancement utilizing parahydrogen. Encyclopedia of Nuclear Magnetic Resonance 9, 750-770 Eds. D. M. Grant and R. K. Harris (2002).
- [0106] 17. Canet, D. et al. Parahydrogen enrichment and hyperpolarization. *Concepts in Magnetic Resonance Part A* 28A, 321-330 (2006).
- [0107] 18. Adams et al. Reversible interactions with parahydrogen enhance NMR sensitivity by polarization transfer, *Science* 323, 1708-1711 (2009).
- [0108] 19. Atkinson, K. D. et al., Spontaneous transfer of parahydrogen induced spin order to pyridine at low magnetic field, *J. Am. Chem. Soc.* 131, 13362-13368 (2009).
- [0109] 20. Chapovsky, P. L. et al. Separation and conversion of nuclear spin isomers of ethylene. *Chem. Phys. Lett.* 322, 424-428 (2000).
- [0110] 21. Sun, Z.-D., Takagi, K., Matsushima, F. Separation and conversion of four nuclear spin isomers of ethylene. *Science* 310, 1938-1941 (2005).
- [0111] 22. Aime, S., Gobetto, R., Reineri, F., Canet, D., Polarization transfer from parahydrogen to heteronuclei: The effect of H/D substitution. The case of the AA'X and A₂A'₂X spin systems. *J. Magn. Res.*, 178, 184-192 (2006).
- [0112] 23. Caravetta, M., Johannessen, O. G. & Levitt, M. H. Beyond the T₁ limit: singlet nuclear spin states in low magnetic fields. *Phys. Rev. Lett.* 92, 153001-153004 (2004).
- [0113] 24. Pileio, G., Caravetta, M. & Levitt, M. H. Extremely Low-Frequency Spectroscopy in Low-Field Nuclear Magnetic Resonance. *Phys. Rev. Lett.* 103, 083002/1-4 (2009).
- [0114] 25. Zax, D. B., Bielecki, A., Zilm, K. W., & Pines, A. Heteronuclear Zero-Field NMR, *Chem. Phys. Lett.* 106, 550-553 (1984).
- [0115] 26. Schaefer, T., Chan, W. K., Sebastian, R., Schurko, R., & Hruška, F. E. Concerning the internal rotational barrier and the experimental and theoretical ⁿJ(¹³C, ¹³C) and ⁿJ(¹H, ¹³C) in ethylbenzene-β¹³C. *Can. J. Chem.* 72, 1972-1977 (1994).
- [0116] 27. Appelt, S. et al. Paths from weak to strong coupling in NMR, *Phys. Rev. A* 81, 023420/1-11 (2010).
- [0117] 28. Lee, C. J., Suter, D., and Pines, A., Theory of multiple-pulse NMR at low and zero fields, *J. Magn. Res.* 75, 110-124 (1987).
- [0118] 29. Dang, H. B., Maloof, A. C., & Romalis, M. V., Ultrahigh sensitivity magnetic field and magnetization measurements with an atomic magnetometer, *Apl. Phys. Lett.* 97, 151110/1-3 (2010).
- [0119] 30. Koptyug, I. V. et al. Para-hydrogen induced polarization in heterogeneous hydrogenation reactions. *J. Am. Chem. Soc.* 129, 5580-5586 (2007)
- [0120] 31. Osborn, J. A., Jardine, F. H., Young, J. F. & Wilkinson, G. The Preparation and Properties of Tris(triphenylphosphine)halogenorhodium(I) and Some Reactions Thereof Including Catalytic Homogeneous Hydrogenation of Olefins and Acetylenes and Their Derivatives. *J. Chem. Soc. A*, 1711-1732 (1966).

CONCLUSION

- [0121] It is to be understood that the above description and examples are intended to be illustrative and not restrictive. Many embodiments will be apparent to those of skill in the art upon reading the above description and examples. The scope of the invention should, therefore, be determined not with reference to the above description and examples, but should instead be determined with reference to the appended claims, along with the full scope of equivalents to which such claims are entitled. The disclosures of all articles and references, including patents, patent applications, and publications, are incorporated herein by reference for all purposes.